

**B.Sc. (Part—I) (Semester—I) Examination**  
**MATHEMATICS**  
**(Algebra & Trigonometry)**  
**Paper—I**

Time : Three Hours]

[Maximum Marks : 60

- Note :—** (1) Question **ONE** is compulsory. Attempt once.  
 (2) Attempt **ONE** question from each Unit.

1. Choose the correct alternative :—

- (1) Which **one** of the following statements is true :— 10
- (a)  $\cosh(x + iy) = \cosh x \cdot \cos y + i \sinh x \cdot \sin y$   
 (b)  $\cosh(x + iy) = \cos x \cos y + i \sin x \sin y$   
 (c)  $\cosh(x + iy) = \cosh x + \cos y - i \sinh x \cdot \sin y$   
 (d)  $\cosh(x + iy) = \cosh x \sin y + i \sinh x \cos y$
- (2) What is the value of  $\sinh^{-1}x$  :
- (a)  $\log[x + \sqrt{x^2 + 1}]$  (b)  $\log[x + \sqrt{x^2 - 1}]$   
 (c)  $\log[x + \sqrt{1 - x^2}]$  (d) None of these
- (3) The value of  $4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{70} + \tan^{-1} \frac{1}{99}$  is \_\_\_\_\_.
- (a)  $\pi/2$  (b)  $\pi/4$   
 (c)  $\pi/3$  (d)  $\pi$
- (4) Sum of the series  $x - \frac{x^2}{2} + \frac{x^3}{3} - \dots + (-1)^{n+1} \frac{x^n}{n} + \dots$ ;  $-1 < x < 1$  is denoted by \_\_\_\_\_.
- (a)  $\log(1 + x)$  (b)  $\sinh x$   
 (c)  $\cosh x$  (d)  $e^x$
- (5) If  $q = 2 + 2i - j + 4k$  then the norm of  $q$  is \_\_\_\_\_.
- (a)  $-5$  (b)  $5$   
 (c)  $1/5$  (d) None of these
- (6) The inverse of unit quaternion is its \_\_\_\_\_.
- (a) Purely imaginary (b) Purely real  
 (c) Complex conjugate (d) None of these
- (7) If  $\alpha + i\beta$  be the root of quadratic polynomial  $f(x) = 0$  then its another root is \_\_\_\_\_.
- (a)  $\alpha - i\beta$  (b)  $\alpha$   
 (c)  $\beta$  (d) None of these
- (8) If  $\alpha, \beta, \gamma$  are the roots of the equation  $px^3 + qx^2 + rx + s = 0$  then  $\Sigma \alpha$  is \_\_\_\_\_.
- (a)  $\frac{q}{p}$  (b)  $-\frac{q}{p}$   
 (c)  $\frac{r}{p}$  (d)  $\frac{s}{p}$

- (9) If A and B are the non-singular matrices of order n then \_\_\_\_\_.
- (a)  $(AB)^{-1} = AB$  (b)  $(AB)^{-1} = A^{-1} \cdot B^{-1}$   
 (c)  $(AB)^{-1} = B^{-1} \cdot A^{-1}$  (b) None of these
- (10) 'Every square matrix satisfies its own characteristics equation' is the statement of \_\_\_\_\_.
- (a) Lagrange's MVT (b) De-Moivre's theorem  
 (c) Cayley-Hamilton theorem (d) Cauchy's MVT

**UNIT—I**

2. (a) Prove that  $\frac{1 + \sin \theta + i \cos \theta}{1 + \sin \theta - i \cos \theta} = \sin \theta + i \cos \theta$ .

Hence prove that  $\left(1 + \sin \frac{\pi}{5} + i \cos \frac{\pi}{5}\right) + i \left(1 + \sin \frac{\pi}{5} - i \cos \frac{\pi}{5}\right) = 0$ . 5

- (b) If  $\sin(\alpha + i\beta) = x + iy$  then prove that  $\frac{x^2}{\cosh^2 \beta} + \frac{y^2}{\sinh^2 \beta} = 1$  and  $\frac{x^2}{\sin^2 \alpha} - \frac{y^2}{\cos^2 \alpha} = 1$  5

3. (p) Prove that one of the value of : 5  
 $(\cos \theta + i \sin \theta)^n$  is  $(\cos n\theta + i \sin n\theta)$ ; when n is negative integer.  
 (q) Separate real and imaginary parts of  $\tan(x + iy)$ . 5

**UNIT—II**

4. (a) Find the Sum of the series :

$$C = 1 + e^{\sin x} \cdot \cos(\cos x) + \frac{1}{2!} e^{2 \sin x} \cdot \cos(2 \cos x) + e^{3 \sin x} \cdot \frac{1}{3!} \cos(3 \cos x) + \dots$$
 5

- (b) Prove that  $4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{70} + \tan^{-1} \frac{1}{99} = \frac{\pi}{4}$ . 5

5. (p) Find the sum of the series  $\sinh x + \frac{1}{2!} \sinh 2x + \frac{1}{3!} \sinh 3x + \dots$  5

- (q) If  $-\frac{\pi}{4} \leq x \leq \frac{\pi}{4}$  then prove that

$$x = \tan x - \frac{1}{3} \tan^3 x + \frac{1}{5} \tan^5 x + \dots + (-1)^{n-1} \frac{1}{2n-1} \tan^{2n-1} x + \dots$$
 5

**UNIT—III**

6. (a) Prove that for  $p, q \in H$ ,  $N(pq) = N(p) N(q)$  and  $N(q^*) = N(q)$ . 5

- (b) For the quaternion  $q = \cos \frac{\pi}{3} + i \sin \frac{\pi}{3}$  and the input vector  $v = i$ , compute the output vector  $w$  under the action of the operators  $L_q$  and  $L_{q^*}$ . 5

7. (p) Show that the quaternion product need not be commutative. 5

- (q) For any  $p, q \in H$ , show that  $pq = qp$  if and only if  $p$  and  $q$  are parallel. 5

**UNIT—IV**

8. (a) Find the roots of the equation,  $8x^3 + 18x^2 - 27x - 27 = 0$ , if these roots are in geometric progression. 5
- (b) State Descartes rule of sign. Find the nature of the roots of the equation  $2x^7 - x^4 + 4x^3 - 5 = 0$ . 5
9. (p) Prove that in an equation with real coefficients complex roots occur in pairs. 5
- (q) Solve the equation  $x^4 - 2x^3 - 22x^2 + 62x - 15 = 0$ ; given that  $2\sqrt{3}$  is one of the root. 5

**UNIT—V**

10. (a) Show that if  $\lambda$  is the eigen value of a nonsingular matrix A then  $\lambda^{-1}$  is the eigen value of  $A^{-1}$ . 5
- (b) Find the eigen values and the corresponding eigen vector for smallest eigen value of the matrix  $A = \begin{bmatrix} 8 & -8 & -2 \\ 4 & -3 & -2 \\ 3 & -4 & 1 \end{bmatrix}$ . 5
11. (p) Show that the eigen values of any square matrix A and A' are same. 5
- (q) Reduce to canonical form and find the rank of the matrix  $A = \begin{bmatrix} 1 & 1 & -1 & 1 \\ 1 & -1 & 2 & -1 \\ 3 & 1 & 0 & 1 \end{bmatrix}$ . 5

